

Solid Tumour Section

Short Communication

Kidney: ALK-rearranged renal cell carcinoma

Adrian Marino-Enriquez

Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA USA;
admarino@partners.org

Published in Atlas Database: February 2017

Online updated version : <http://AtlasGeneticsOncology.org/Tumors/ALKrenalCellCarclD6279.html>

Printable original version : <http://documents.irevues.inist.fr/bitstream/handle/2042/68916/02-2017-ALKrenalCellCarclD6279.pdf>

DOI: 10.4267/2042/68916

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2018 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

Review on ALK-rearranged renal cell carcinoma, summarizing clinical and genetic data

Keywords

Renal cell carcinoma; ALK; HOOK1; TPM3; STRN; EML4; VCL

Identity

Phylum

Urinary system:Kidney: Renal cell tumors: :ALK-rearranged renal cell carcinoma

Classification

ALK-rearranged renal cell carcinoma is a distinct type of renal cell carcinoma included in the so-called emerging/provisional RCC. The 2013 International Society of Urological Pathology (ISUP) Vancouver Classification of (adult) renal neoplasia identified a category of emerging or provisional new entities. Although these entities appeared to be distinct, these are rare tumors not fully characterized and additional reports will be needed to refine their diagnostic criteria and established clinical outcome.

Clinics and pathology

Disease

ALK-rearranged renal cell carcinoma is a distinct emerging type of RCC that commonly affects

children and young adults. The disease is defined as a RCC harboring ALK gene rearrangements, resulting in oncogenic fusions with a variety of partner genes.

Epidemiology

ALK-rearranged renal cell carcinoma is an uncommon type of RCC. Initially described in children, in which it may be over-represented, further investigations demonstrated that a small proportion of adult RCC belong to this ALK-rearranged category. Of 18 well-documented cases reported to date, 8 have been described in children (6-16 yo), and 10 in adults (33-61 yo). An association with sickle cell trait was observed in the first cases described, but this association seems inconsistent and may be limited to cases with a specific VCL/ALK gene fusion.

Clinics

These form masses in the kidney. The majority of cases described so far have been confined to the kidney with a median size of 4.5 cm. Nodal extension at presentation has been observed; distant metastases are rare and have been observed more commonly in adult patients. The clinical course of ALK-rearranged RCC is frequently indolent, although rare cases, more frequently in adult patients, pursue a more aggressive clinical course.

Pathology

Grossly, ALK-rearranged RCC are brown to tan solid masses that may be well circumscribed or

infiltrative, usually located centrally with extension into the renal pelvis. Histologically, the tumors most commonly feature a solid architecture, with occasional trabecular and tubular features, and are composed of sheets of epithelioid cells with abundant pale eosinophilic cytoplasm and frequent intracytoplasmic lumina. Cell nuclei show high-grade features, with vesicular chromatin and prominent nucleoli -usually ISUP grade 3, but occasionally ISUP grade 4. Rhabdoid morphology or intracytoplasmic inclusions may be focally present. There is a prominent capillary network. Less frequently, ALK-rearranged RCC may show a predominantly papillary architecture, mucin deposition and focal psammomatous calcification. It

is unclear if some of these features correlate with specific fusion partners. Immunohistochemically, the tumor cells consistently express ALK, CK7, EMA, and TFE3 (which should not be interpreted as evidence of TFE3 rearrangement, absent in this tumor type); nuclear INI1 expression is retained.

Treatment

Treatment: surgical excision. Treatment with ALK inhibitors has provided clinical benefit in cases with advanced disease.

Cytogenetics

Cytogenetics Morphological

ALK Gene Fusions in renal cell carcinoma (RCC)

Neoplasm	Fusion	Age Range (years) (n)
ALK-rearranged RCC	t(2;10)(p23;q22) VCL/ALK	6-16 (3)
ALK-rearranged RCC	t(1;2)(q25;p23) TPM3/ALK	12-49 (7)
ALK-rearranged RCC	inv(2)(p22p23) STRN/ALK	33,38 (2)
ALK-rearranged RCC	inv(2)(p21p23) EML4/ALK	52,53 (2)
ALK-rearranged RCC	t(1;2)(p32;p23) HOOK1/ALK	16 (1)
ALK-rearranged RCC	ALK/unknown	44-61 (3)

Result of the chromosomal anomaly

Fusion Protein

Description

ALK-rearranged RCC are characterized by fusion of the ALK tyrosine kinase gene with one of several gene partners including VCL, TPM3, STRN, EML4, and HOOK1 (Table 1). The originally described translocation in ALK-rearranged RCC was t(2;10)(p23;q22) which fuses the 5' end of VCL to the 3' end of ALK. So far, VCL/ALK fusions have been described only in ALK-rearranged RCC - interestingly, in 3 tumors affecting pediatric patients with sickle cell trait. Similarly, HOOK1/ALK fusion has been identified only in RCC. The TPM3/ALK, STRN/ALK and EML4/ALK gene fusions, however, are similar to those found in other tumor types such as inflammatory myofibroblastic sarcoma, thyroid carcinoma and lung adenocarcinoma. All of these ALK gene fusions result in oncogenic proteins that include the kinase domain of ALK, fused to structural protein motifs

that enable direct or indirect oligomerization. The ALK fusion partners also contribute active promoters that lead to strong expression of ALK, which can be detected by immunohistochemistry.

References

- Cajal MM, Jennings LJ, George D, Perlman EJ. Expanding the spectrum of ALK-rearranged renal cell carcinomas in children: Identification of a novel HOOK1-ALK fusion transcript. *Genes Chromosomes Cancer*. 2016 Oct;55(10):814-7
- Chen YB, Xu J, Skanderup AJ, Dong Y, Brannon AR, Wang L, Won HH, Wang PI, Nanjangud GJ, Jungbluth AA, Li W, Ojeda V, Hakimi AA, Voss MH, Schultz N, Motzer RJ, Russo P, Cheng EH, Giannotti FG, Lee W, Berger MF, Tickoo SK, Reuter VE, Hsieh JJ. Molecular analysis of aggressive renal cell carcinoma with unclassified histology reveals distinct subsets. *Nat Commun*. 2016 Oct 7;7:13131
- Debelenko LV, Raimondi SC, Daw N, Shivakumar BR, Huang D, Nelson M, Bridge JA. Renal cell carcinoma with novel VCL-ALK fusion: new representative of ALK-associated tumor spectrum. *Mod Pathol*. 2011 Mar;24(3):430-42
- Jeanneau M, Gregoire V, Desplechain C, Escande F, Tica DP, Aubert S, Leroy X. ALK rearrangements-associated

Kidney: ALK-rearranged renal cell carcinoma

renal cell carcinoma (RCC) with unique pathological features in an adult. *Pathol Res Pract*. 2016 Nov;212(11):1064-1066

Kusano H, Togashi Y, Akiba J, Moriya F, Baba K, Matsuzaki N, Yuba Y, Shiraishi Y, Kanamaru H, Kuroda N, Sakata S, Takeuchi K, Yano H. Two Cases of Renal Cell Carcinoma Harboring a Novel STRN-ALK Fusion Gene. *Am J Surg Pathol*. 2016 Jun;40(6):761-9

Lee C, Park JW, Suh JH, Nam KH, Moon KC. ALK-Positive Renal Cell Carcinoma in a Large Series of Consecutively Resected Korean Renal Cell Carcinoma Patients. *Korean J Pathol*. 2013 Oct;47(5):452-7

Mariño-Enríquez A, Ou WB, Weldon CB, Fletcher JA, Pérez-Atayde AR. ALK rearrangement in sickle cell trait-associated renal medullary carcinoma. *Genes Chromosomes Cancer*. 2011 Mar;50(3):146-53

Moch, H., Humphrey, P.A., Ulbright, T.M., Reuter, V.E. eds (4th edition);. *World Health Organization Classification of Tumours of the Urinary Systems and Male Genital Organs* IARC Press, Lyon 2016; 16-17.

Smith NE, Deyrup AT, Mariño-Enríquez A, Fletcher JA, Bridge JA, Illei PB, Netto GJ, Argani P.. VCL-ALK renal cell carcinoma in children with sickle-cell trait: the eighth sickle-cell nephropathy? *Am J Surg Pathol*. 2014 Jun;38(6):858-63.

Srigley JR, Delahunt B, Eble JN, Egevad L, Epstein JI, Grignon D, Hes O, Moch H, Montironi R, Tickoo SK, Zhou M, Argani P; ISUP Renal Tumor Panel.. The International Society of Urological Pathology (ISUP) Vancouver Classification of Renal Neoplasia.

Sukov WR, Hodge JC, Lohse CM, Akre MK, Leibovich BC, Thompson RH, Cheville JC.. ALK alterations in adult renal cell carcinoma: frequency, clinicopathologic features and outcome in a large series of consecutively treated patients. *Mod Pathol*. 2012 Nov;25(11):1516-25.

Thorner PS, Shago M, Marrano P, Shaikh F, Somers GR.. TFE3-positive renal cell carcinomas are not always Xp11 translocation carcinomas: Report of a case with a TPM3-ALK translocation. *Pathol Res Pract*. 2016 Oct;212(10):937-942.

Yu W, Wang Y, Jiang Y, Zhang W, Li Y.. Genetic analysis and clinicopathological features of ALK-rearranged renal cell carcinoma in a large series of resected Chinese renal cell carcinoma patients and literature review. *Histopathology*. 2017 Feb 15. doi: 10.1111/his.13185.

This article should be referenced as such:

Marino-Enriquez A. Kidney: ALK-rearranged renal cell carcinoma. *Atlas Genet Cytogenet Oncol Haematol*. 2018; 22(6):260-262.
